

EW PORTAL TO THE COMPOSITE DM

HIDDEN VALLEY

Lingfeng Li
李凌风

Nov. 07, 2026

河南师范大学
HENAN NORMAL UNIVERSITY




26xx.abcd H-C. Cheng, X. Jiang, LFL, E. Salvioni

2408.13304 H-C. Cheng, X. Jiang, LFL

2401.08785 H-C. Cheng, X. Jiang, LFL, E. Salvioni

2110.10691 H-C. Cheng, LFL, E. Salvioni



The overview of this talk

The Electroweak Portal...

- The legacy of SM as a tool
- “Irrelevant” portal EFT
- EW precision tests
- Flavor portal

...to the Hidden Valley ...

- Dark confinement
- Long-lived particles
- Dark hadrons & symmetry
- Phenomenology

...and Beyond

- Composite Dark Matter
- Cosmological Probes


COSMIC


RICHNESS

GENERIC


Higgs

EW






**CMB Power
Spectra**




**Large Scale
Structure**




**Galaxy
Formation**



Lensing



Composite DM Candidates





Dark Baryons



Dark R-hadrons


Dark Mesons



Dark Blueballs

Motivation

"Baryons" are Composite




Hierarchy
Asymmetry

GUT


FOPT

CFT




UV Model Building

"Minimal" Setup (?)




G
only?



Unique Dynamics & Phenomena

- Planetary science implications
- New phys. from extremes conditions



J. Fan, LFL, JHEP 10, 186 (2022); J. Fan, LFL, C. Sun, 2501.12440; S. Yan, LFL, J. Fan, JHEP 06, 028 (2024), Phys. Rev. Lett. 135, 231801 (2025)

S. Chai, J. Gu, LFL JHEP 05, 292 (2024); T.S. M. Ho, X. H. Jiang, T. H. Kwok, LFL, T. Liu, Phys. Rev. D 109, no.9, 093004 (2024);

- Pheno of Hidden Valley
- Flavor tests of SM
- Machine learning developments

- Primordial features & non-Gaussianity
- Ultralight DM and axion physics



LFL, S. Lu, Y. Wang, S. Zhou, JHEP 07, 231 (2020); L. Fong, LDL, T. Liu, H. Luu, T. Qiu, H. Tye, JCAP 08, 057 (2021); Y. Bao, J. Fan, LFL, Phys. Rev. Lett. 130, no.24, 241001 (2023); X. Chen, J. Fan, LFL, JHEP 12, 197 (2023)

SM Legacy as Tools

- SM is “complete” and known with reasonable precision


Standard Model Total Production Cross Section Measurements

Status: October 2023



- What can we learn from all these?

The Irrelevant Portal



$\epsilon \mathcal{O}_{\text{SM}} J_{\text{Dark}}$ **Relevant portal:** small couplings to keep the valley hidden

$\frac{1}{\Lambda^n} \mathcal{O}_{\text{SM}} J_{\text{Dark}}$ **Irrelevant portal:** Hidden behind EFT


The Electroweak Portal

$$\frac{c}{M^2} (iH^\dagger \overleftrightarrow{D}_\mu H) (\bar{\psi} \gamma^\mu \psi)$$



h

Z




EW bosons stand for a scale of 246 GeV, close enough to TeV+

>TeV Scale $\approx$ Dark Enough

The EW Portal: $\frac{c}{M^2} (iH^\dagger \overleftrightarrow{D}_\mu H) (\bar{\psi} \gamma^\mu \psi)$ from Two UV Setups


$$\mathcal{L}_{\text{UV}} \supset \overline{Q} Y \psi H \quad \longrightarrow \quad \frac{c}{M^2} \sim \frac{Y^2}{M_Q^2}$$



Heavy fermion doublet mediation

H-C. Cheng, LFL, E.
Salvioni, 2110.10691

$$\mathcal{L}_{\text{UV}} \supset \delta \hat{M}^2 Z^\mu Z'_\mu \xrightarrow{c} \frac{c}{M^2} \sim \frac{g_D^2 \delta \hat{M}^2}{m_Z^2 m_{Z'}^2}$$



Dark Z' mediation: a Z' heavier or lighter(!) than Z with mass mixing

H-C. Cheng, X. Jiang, LFL, E. Salvioni, 2401.08785

See also Y. C. San , M. Perelstein, and P. Tanedo, 2022

Exotic BR of Z

A key phenomenon for EW portal models

As the doublet mediation benchmark:

$$\text{BR}(Z \rightarrow \psi' \bar{\psi}') \approx 1.8 \times 10^{-4} \left(\frac{N_d \text{Tr}(\mathbf{Y} \mathbf{Y}^\dagger \mathbf{Y} \mathbf{Y}^\dagger)}{3} \right) \left(\frac{1 \text{ TeV}}{M} \right)^4$$

For the Z' mediation :

$$\text{BR}(Z \rightarrow \psi' \bar{\psi}') \approx 1.4 \times 10^{-4} \left(\frac{N_d}{3} \right) \left(\frac{\xi}{0.01} \right)^2 g_D^2 \sum_i (x_{Li}^2 + x_{Ri}^2)$$

$$\text{The effective mixing angle } \xi \sim \frac{\delta M^2}{M_Z^2 - M_{Z'}^2}$$


Exotic BR of h is not large ($< \sim 10^{-4}$), the constraints are weaker than those from Z decays ($\sim 10^3$ more Z than h at LHC)

EWPT Corrections

M.E. Peskin, T. Takeuchi, 1990

M.E. Peskin, T. Takeuchi, 1992

Doublet mediation



Leading by EW loops with mixing

L. Lavoura, J. P. Silva, 1993;

C. Anastasiou, E. Furlan, J.

Santiago, 2009

$$S \simeq \frac{2N_d}{45\pi M^2} \left(\frac{5Y^2 v^2}{2} - 3m_Z^2(1 - 2s_W^2 + 2s_W^4) \right)$$

$$T \simeq \frac{N_d y_{11}^4 v^4}{48\pi s_W^2 m_W^2 M^2}$$

Z' mediation



Leading by LEP Z pole observables defined with:

$$\mathcal{L}_{Zf\bar{f}} = -\frac{\bar{Z}e}{s_W c_W} \bar{f} \gamma^\mu (T_{Lf}^3 - s_*^2 Q_f) f Z_\mu$$

B. Holdom, 1991; G. Altarelli

et. al, 1991

$$S = \frac{4s_W^2}{\alpha} \left(\frac{c_W^2}{s_W} \xi t_\chi - c_W^2 \xi^2 \right)$$



$$T = \frac{1}{\alpha} \left(2s_W \xi t_\chi + \xi^2 \left(\frac{M_{Z'}^2}{M_Z^2} - 2 \right) \right)$$

The Flavor Portal

$$\frac{c}{M^2} (iH^\dagger \overleftrightarrow{D}_\mu H) (\bar{\psi} \gamma^\mu \psi) \quad \text{Tree level, meson decay and dark shower}$$



$h \ Z$



$$\frac{\mathcal{K}}{M^2} (\bar{d}_i \gamma^\mu d_j) (\bar{\psi} \gamma_\mu \psi)$$

W. Altmannshofer, A. Crivellin, H. Haigh, G. Inguglia and J. Martin Camalich, 2311.14629; J. Martin Camalich, M. Pospelov, P. N. H. Vuong, R. Ziegler and J. Zupan, 2002.04623; M. K. Gaillard, M. B. Gavela, R. Houtz, P. Quilez and R. Del Rey, 1805.06465; T. Li, M.A. Schmidt and M. Yuan, 2506.08342.....


Flavor Portal from EW only

$\Gamma_{B,K,\dots}$ are suppressed by $(M_W)^{-4}$ in SM.

Inami, Lim 1980

X.G. He, J. Tandean,

G. Valencia, 0909.3638



The four-fermion EFT respecting MFV


$$-\frac{c}{M^2} \frac{g^2}{128\pi^2} \underline{J_D^\mu \bar{d}_j \gamma_\mu P_L d_i} \sum_{q \in u,c,t} \underline{V_{qj}^* V_{qi} \mathcal{K}_q}$$

Flavor Portal from EW only

Inami, Lim 1980

X.G. He, J. Tandean,

G. Valencia, 0909.3638



In heavy doublet mediation:

$$\mathcal{K}_q \equiv 3 + x_q \left(\log \frac{\Lambda_{UV}^2}{m_q^2} - 2 \right) \quad x_q \equiv m_q^2 / M_W^2$$

In Z' mediation (the mZ' terms cancel out):

Typically $O(10)$

$$\mathcal{K}_q \equiv x_q \log \frac{\Lambda_{UV}^2}{M_W^2} + \frac{-7x_q + x_q^2}{2(1 - x_q)} - \frac{4x_q - 2x_q^2 + x_q^3}{(1 - x_q)^2} \log x_q$$

Does it constrain anything?


We can barely see the SM Z & W in $bs\nu\nu$ and $sd\nu\nu$
(for now...)

Γ_{39}	$\pi^+ \nu \bar{\nu}$	$(1.14^{+0.40}_{-0.33}) \times 10^{-10}$
Γ_{40}	$\pi^+ \pi^0 \nu \bar{\nu}$	$< 4.3 \times 10^{-5}$
Γ_{582}	$\pi^+ \nu \bar{\nu}$	$< 1.4 \times 10^{-5}$
Γ_{588}	$K^+ \bar{\nu} \nu$	$< 1.6 \times 10^{-5}$
Γ_{589}	$\rho^+ \nu \bar{\nu}$	$< 3.0 \times 10^{-5}$

Want to Kick Things Up?




Hidden Valley Models



M. J. Strassler and K. M. Zurek, 0604261

$$\Gamma \sim \text{Tr}[Y^\dagger Y \sigma_{\text{hadron}}] \text{ or } \text{Tr}[X \sigma_{\text{hadron}}]$$


Symmetries Yes



The **Dark Force** for Confinement

➤ Neutral naturalness calls
for a non-QCD color

Z. Chacko, H.S. Goh, and R. Harnik, 0506256;
G. Burdman, Z. Chacko, H.S. Goh and
R. Harnik, 0609152;
H-C. Cheng, LL, E. Salvioni, and
C. Verhaaren, 1803.03561




➤ Makes good dark
matter candidate



Y. Hochberg, E. Kuflik,
H. Murayama, T. Volansky,
J. Wacker, 1411.3727;
A. Katz, E. Salvioni, and
B. B. Shakya, 2006.15148; s


P. Schwaller, D. Stolarski, A.
Weiler, 1502.05409;
CMS, 1810.10069;
S. Knapen, J. Shelton, D. Xu,
2103.01238;
S. Born, R. Karur, S. Knapen,
J. Shelton, 2303.04167;
J. Carroasco, J. Zivita, 2307.04847



➤ Rich LLP
phenomenology


➤ Duality ? ALPs ? Baryogenesis?

The Cartoon of Dark Spectrum



Dark Pions

Z portal dark pion production



Dark pions rearrange into **CP eigenstates** (like K_S and K_L in the SM)

The π_1 and π_3 decay via Z portal, ALP-like (axion-like-particle) with **effective ALP decay constants**:


$$\mathcal{L}_a \supset -\frac{\partial_\mu a}{f_a} \sum_f T_f^3 \bar{f} \gamma^\mu \gamma^5 f$$

$$f_a \propto \frac{M^2}{f_{\hat{\pi}}} \quad \text{Just a trick!}$$

Z portal dark pion decay


The π_2 mix with the Higgs since it's CP-even, with **mixing angle**:

$$\mathcal{L}_s \supset -\sin \theta_s \frac{m_f}{v} s \bar{f} f, \quad \theta_s \lesssim 10^{-6}$$



Dark Pions

Z portal dark pion production



Dark pions rearrange into **CP eigenstates** (like K_S and K_L in the SM)

The π_1 and π_3 decay via Z portal, ALP-like (axion-like-particle) with **effective ALP decay constants**:

$$\mathcal{L}_a \supset -\frac{\partial_\mu a}{f_a} \sum_f T_f^3 \bar{f} \gamma^\mu \gamma^5 f$$

$$f_a \propto \frac{M^2}{f_{\hat{\pi}}} \quad \text{Just a trick!}$$

$f_{\hat{\pi}}$ The dark meson decay constant, O(confinement scale)

f_a The effective (composite) ALP decay constant, O(PeV) or above


Convenient for the matching with existing ALP searches

Dark (Flavor) Symmetry

- Dark isospin symmetry: identical mass and couplings $\Rightarrow$ All dark pions stable, DM?
- Dark U(1) symmetry: mass and charge matrix diagonal $\Rightarrow$ dark pions carrying non-zero U(1) global charge are stable, DM?
- Dark CP-symmetry: all entries above are real $\Rightarrow$ No mixing between CP-odd and –even species
-

Dark Pion Decays (ALP-Like)

Recent Update: [R. Balkin, T. Coren, Y. Soreq and M. Williams, 2506.15637](#)



Decay at Human/Lab Length Scales



See also:

[D. Aloni, Y. Soreq and M. Williams, 1811.03474](#); [M. Ovchinnikov and A.~Zaporozhchenko, 2501.04525](#); [Y. Bai, T. K. Chen, J. Liu and X. Ma, 2505.24822](#)

Long-Lived Particles




They are more common and fundamental than it appears



..... neutral
 ——— charged
 ——— any charge

■ BSM
 ■ lepton
 ■ quark
 ■ photon
 ■ anything



P. Schwaller, D. Stolarski, A. Weiler, 1502.05409;
 CMS, 1810.10069;
 S. Knapen, J. Shelton, D. Xu, 2103.01238;
 S. Born, R. Karur, S. Knapen, J. Shelton, 2303.04167;
 J. Carroasco, J. Zircita, 2307.04847


Dark Shower & Emerging Jets

If any dark pion is a LLP $\rightarrow$ emerging energy outside the innermost detector


M. J. Strassler and K. M. Zurek, 0604261

P. Schwaller, D. Stolarski and A. Weiler, 1502.05409

T. Cohen, M. Lisanti and H. K. Lou, 1503.00009





Flavor Portal to LLP



➤ 5-10 times larger BR for inclusive X_s decays


$$\text{BR}(B^{+,0} \rightarrow \{K^+ \hat{\pi}, K^{*0} \hat{\pi}\}) \approx \{0.92, 1.1\} \times 10^{-8} \left(\frac{1 \text{ PeV}}{f_a} \right)^2 \left(\frac{\mathcal{K}_t}{10} \right)^2 \{ \lambda_{BK\hat{\pi}}^{1/2}, \lambda_{BK^*\hat{\pi}}^{3/2} \}$$

$$\text{BR}(K^+ \rightarrow \pi^+ \hat{\pi}) \approx 3.9 \times 10^{-11} \left(\frac{1 \text{ PeV}}{f_a} \right)^2 \left(\frac{\mathcal{K}_t}{10} \right)^2 \lambda_{K\pi\hat{\pi}}^{1/2}$$




LLP Phenomenology

Dark Shower




Signal: Muon rich jet-like structure with long-lived tracks + MET

LHCb




CMS




Searching for displaced dimuon resonances (one or more)

[LHCb, 2007.03923](#); [CMS, 2112.13769](#)




$$BR \approx 10^{-8} \left(\frac{1 \text{ PeV}}{f_a} \right)^2$$




FCNC Decays

LLP at LHC Detectors




LHCb Vertex Locator (VELO)





LHCb: “Trigger-free” Searches


See [LHCb, 2007.03923](#)



Matching with multiple ways to calibrate the projection


$$\text{BR}(Z \rightarrow \bar{\psi}\psi) \times \text{BR}(\hat{\pi} \rightarrow \mu^+ \mu^-)$$






➤ Extended to the HL-LHC era, improve by $\sim \text{O}(10^2)$

➤ Also sensitive to FCNC, unsuppressed B production in the forward region





LHCb: Benchmark




Higgs portal results not
shown here

CMS: Dimuon Scouting

CMS, 2112.13769




- Drop-out detailed info to boost dimuon trigger rate, useful for low mass resonances
- Not bkg free




- Applies to Dark shower or FCNC

CMS: Model-Independent Limits


- Current limit with <11 cm transverse displacement
- 30 times luminosity, detector/algorithm upgrades.....
- ↘ HL-LHC limits









- At HL-LHC, high efficiency to do 2DV search: two dimuon DV with the same mass




Auxiliary Detector Targeting LLPs




MATHS

The logo for FASER, featuring the word "FASER" in a bold, italicized, sans-serif font, with a stylized swoosh or arrow graphic underneath it.


Auxiliary LLP Detectors







Corresponding f_a range increase up to $O(100)$ PeV

Rare/Precision/Intensity Frontier







Prompt Dark Vector from DS



- Exotic Higgs or Z decays
- Recast from ATLAS $Z \rightarrow 4(6)$ lepton exotic searches

ATLAS 2306.07413




J. Cheng, R. Husain,
 LFL, M. Strassler,
 2412.14452 + in prep.

Time to Find out THE Dark but Colorful



GENERIC✓
RICHNESS✓

Back in the SM



Thermal Mechanisms



The messenger: heavy meson that decays, contact with the SM



The DM meson, non-interacting



The intermediate, exists in many models



SIMP-like



Pairwise



Scattering



Y. Hochberg, E.
Kuflik, H.
Murayama, T.
Volansky, J.
Wacker, 1411.3727;
A. Katz, E.
Salvioni, and B.
Shakya,
2006.15148.....

The Model

Gauge group: $SO(N_c)$ $N_c \geq 4$

$SU(3)/SO(3) \Rightarrow$ 5 Goldstone bosons

Two π , two K and one η




$$m_\eta = (1 + \Delta_\eta) m_\pi > m_K = (1 + \Delta_K) m_\pi > m_\pi$$

Leading ChPT Lagrangian (Low Temp)

$$\begin{aligned}
\mathcal{L}_2 \supset & \frac{1}{24f^2} \left\{ m_{\hat{\pi}}^2 \left[\hat{\eta}_0^2 + 2\hat{\pi}_+\hat{\pi}_- + 2\hat{K}_{+1/2}\hat{K}_{-1/2} \right]^2 + 2(m_{\hat{K}}^2 - m_{\hat{\pi}}^2) \left[\frac{8}{9}\hat{\eta}_0^4 + 3\hat{\eta}_0^2\hat{K}_{+1/2}\hat{K}_{-1/2} \right. \right. \\
& - \left. \left. \sqrt{\frac{2}{3}}(\hat{\eta}_0\hat{K}_{+1/2}^2\hat{\pi}_- + \hat{\eta}_0\hat{K}_{-1/2}^2\hat{\pi}_+) + 2\hat{K}_{+1/2}^2\hat{K}_{-1/2}^2 + 2\hat{K}_{+1/2}\hat{K}_{-1/2}\hat{\pi}_+\hat{\pi}_- \right] \right\} \\
& + \frac{1}{24f^2} \left\{ 4(\partial\hat{\pi}_+)^2\hat{\pi}_-^2 + 4\hat{\pi}_+^2(\partial\hat{\pi}_-)^2 - 8\partial\hat{\pi}_+\hat{\pi}_+\partial\hat{\pi}_-\hat{\pi}_- + (\partial\hat{K}_{+1/2})^2\hat{K}_{-1/2}^2 + \hat{K}_{+1/2}^2(\partial\hat{K}_{-1/2})^2 \right. \\
& - 2\partial\hat{K}_{+1/2}\hat{K}_{+1/2}\partial\hat{K}_{-1/2}\hat{K}_{-1/2} + 8\hat{K}_{+1/2}\partial\hat{K}_{-1/2}\hat{\pi}_+\partial\hat{\pi}_- + 8\partial\hat{K}_{+1/2}\hat{K}_{-1/2}\partial\hat{\pi}_+\hat{\pi}_- \\
& - 4\hat{K}_{+1/2}\hat{K}_{-1/2}\partial\hat{\pi}_+\partial\hat{\pi}_- - 4\hat{K}_{+1/2}\partial\hat{K}_{-1/2}\partial\hat{\pi}_+\hat{\pi}_- - 4\partial\hat{K}_{+1/2}\hat{K}_{-1/2}\hat{\pi}_+\partial\hat{\pi}_- \\
& - 4\partial\hat{K}_{+1/2}\partial\hat{K}_{-1/2}\hat{\pi}_+\hat{\pi}_- - 6\hat{\eta}_0^2\partial\hat{K}_{+1/2}\partial\hat{K}_{-1/2} - 6(\partial\hat{\eta}_0)^2\hat{K}_{+1/2}\hat{K}_{-1/2} \\
& + 6\hat{\eta}_0\partial\hat{\eta}_0\partial\hat{K}_{+1/2}\hat{K}_{-1/2} + 6\hat{\eta}_0\partial\hat{\eta}_0\hat{K}_{+1/2}\partial\hat{K}_{-1/2} + 2\sqrt{6} \left[\partial\hat{\eta}_0\partial\hat{K}_{-1/2}\hat{K}_{-1/2}\hat{\pi}_+ \right. \\
& + \partial\hat{\eta}_0\partial\hat{K}_{+1/2}\hat{K}_{+1/2}\hat{\pi}_- - \partial\hat{\eta}_0\hat{K}_{+1/2}^2\partial\hat{\pi}_- - \partial\hat{\eta}_0\hat{K}_{-1/2}^2\partial\hat{\pi}_+ - \hat{\eta}_0(\partial\hat{K}_{-1/2})^2\hat{\pi}_+ \\
& \left. - \hat{\eta}_0(\partial\hat{K}_{+1/2})^2\hat{\pi}_- + \hat{\eta}_0\partial\hat{K}_{-1/2}\hat{K}_{-1/2}\partial\hat{\pi}_+ + \hat{\eta}_0\partial\hat{K}_{+1/2}\hat{K}_{+1/2}\partial\hat{\pi}_- \right] \left. \right\}.
\end{aligned}$$

**Mass & 4-point
interaction terms**


Forbidden Composite DM



**Only energetic particle can go through
the mass gap, exponential sensitivity**

R.T. D'Agnolo and J.T. Ruderman, 1505.07107; LFL, Y. Tsai, 1901.0993; E. Bernreuther, F. Kahlhoefer, M. Kramer and P. Tunney, 1907.04346.....


Important Alternative Processes




K assisted conversion, low Boltzmann suppression

K annihilation, gives indirect detection signals


Large Decay Width Limit: Freezeout Controlled by Conversion



Small Decay Width Limit: Freezeout Controlled by the Decay




Decay at
Human/Lab
Length Scales



Extended Phases


At least 3 extra dynamic phases recognized

* Assuming all processes having similar cross sections, not necessarily true for realistic models





Examining Conversion Rates:


No conversion limit,
K as secondary DM



Full $\pi K \rightarrow \eta K$
dominance








Direct Detection Prospects

The leading Z exchange between DM and SM are forbidden by the same symmetry that stabilizes the DM, the remaining Higgs exchange:

$$\mathcal{L} \supset 4B \frac{y_1 \tilde{y}_1}{M_{Q_1}} \pi_+ \pi_- |H|^2 ,$$




Nucleon
scattering $\sigma_{\text{see}} \approx 2.5 \times 10^{-45} \text{ cm}^2 b^2 \left(\frac{m_N}{m_\pi + m_N} \right)^2 \left(\frac{f}{\text{GeV}} \right)^2 \left(\frac{y_1 \tilde{y}_1}{10^{-2}} \right)^2 \left(\frac{\text{TeV}}{M_{Q_1}} \right)^2$

When $m > 8 \text{ GeV}$, could be above the neutrino fog

Indirect Detection Prospects

- As forbidden dark matter, usually free from indirect detection
- The remaining secondary Kaon DM can annihilate to SM, strongly depending on how many K remain




The most stringent CMB spectral distortion limit
constraints the highly degenerate case ($\Delta < 1\%$)

Recent progress: freeze-in of a composite
dark sector LFL, Sida Lu, In preparation

Summary

- EW portal to the dark is a generic feature to be searched
- Dark confinement gives rise to LLP with good motivations
- At GeV scale, current and future limits are projected
- May lead to composite DM and non-trivial cosmology



Backup Slides

Lagrangian ($\mathbf{Z'}$ Mediation)

$$\mathcal{L}_{\text{mix}} = -\frac{\sin \chi}{2} \hat{Z}'_{\mu\nu} \hat{B}^{\mu\nu} + \delta \hat{M}^2 \hat{Z}'^\mu \hat{Z}_\mu - \kappa \Phi^* \Phi H^\dagger H$$

$$\begin{pmatrix} A_\mu \\ Z_\mu \\ Z'_\mu \end{pmatrix} = L^{-1} \begin{pmatrix} \hat{A}_\mu \\ \hat{Z}_\mu \\ \hat{Z}'_\mu \end{pmatrix} \quad L = \begin{pmatrix} 1 & -\hat{c}_W \sin \xi \tan \chi & -\hat{c}_W \cos \xi \tan \chi \\ 0 & \cos \xi + \hat{s}_W \sin \xi \tan \chi & -\sin \xi + \hat{s}_W \cos \xi \tan \chi \\ 0 & \sin \xi / \cos \chi & \cos \xi / \cos \chi \end{pmatrix}$$

$$\tan 2\xi = \frac{-2 \cos \chi (\delta \hat{M}^2 + \hat{M}_Z^2 \hat{s}_W \sin \chi)}{\hat{M}_{Z'}^2 - \hat{M}_Z^2 \cos^2 \chi + \hat{M}_Z^2 \hat{s}_W^2 \sin^2 \chi + 2\delta \hat{M}^2 \hat{s}_W \sin \chi}$$

Lagrangian (Fermion Mediation)

$$- \mathcal{L}_{\text{UV}} = \overline{Q}_L \mathbf{Y} \psi_R H + \overline{Q}_R \tilde{\mathbf{Y}} \psi_L H + \overline{Q}_L \mathbf{M} Q_R + \overline{\psi}_L \boldsymbol{\omega} \psi_R + \text{h.c.}$$

$$\begin{aligned} \mathcal{L}_{\text{EFT}} = & \frac{1}{2} \overline{\psi}_R \mathbf{Y}^\dagger \mathbf{M}^{-2} \mathbf{Y} \left[|H|^2 i \not{D} + i \gamma^\mu H^\dagger D_\mu H \right] \psi_R + \text{h.c.} \\ & + \frac{1}{2} \overline{\psi}_L \tilde{\mathbf{Y}}^\dagger \mathbf{M}^{-2} \tilde{\mathbf{Y}} \left[|H|^2 i \not{D} + i \gamma^\mu H^\dagger D_\mu H \right] \psi_L + \text{h.c.} \\ & - \overline{\psi}_L \boldsymbol{\omega} \psi_R + \overline{\psi}_L \tilde{\mathbf{Y}}^\dagger \mathbf{M}^{-1} \mathbf{Y} \psi_R |H|^2 + \text{h.c.} , \end{aligned}$$

Z Pole EWPT (Z' Mediation)

$$\mathcal{L}_{Zf\bar{f}} = -\frac{\bar{Z}e}{s_W c_W} \bar{f} \gamma^\mu (T_{Lf}^3 - s_*^2 Q_f) f Z_\mu$$

$$\bar{Z} = 1 + \frac{\alpha T}{2}, \quad s_*^2 = s_W^2 + \frac{\alpha}{c_W^2 - s_W^2} \left(\frac{S}{4} - s_W^2 c_W^2 T \right)$$

$$S = \frac{4s_W^2}{\alpha} \left(\frac{c_W^2}{s_W} \xi t_\chi - c_W^2 \xi^2 \right)$$

$$T = \frac{1}{\alpha} \left(2s_W \xi t_\chi + \xi^2 \left(\frac{M_{Z'}^2}{M_Z^2} - 2 \right) \right)$$

$$U = \frac{4s_W^2}{\alpha} c_W^2 \xi^2$$

Flashback I: Accidental SUSY with Neutral Naturalness

$$W'_{Z_3} = y_t(Q_A H u_A^c + Q_B H u_B^c + Q_C H u_C^c) + \omega(u'_B u_B^c + u'_C u_C^c) + M(Q_B Q_B'^c + Q_C Q_C'^c)$$


$$V'_s = \tilde{m}^2(|\tilde{Q}_A|^2 + |\tilde{u}_A^c|^2) - \tilde{m}^2(|\tilde{Q}_B|^2 + |\tilde{Q}_C|^2)$$

$Q_A \ u_A^c$ SM top sector

$$Q_{B,C} = \begin{pmatrix} t_{B,C} \\ b_{B,C} \end{pmatrix} \sim 2_{-1/2}$$


$$u_{B/C}^c \sim 1_0$$

- Only compatible with the LHC if dark color exists



Flashback II: Composite DM

$$m_{\eta_d} = 10, m_{\omega_d} = 12 \text{ GeV}, c\tau(\omega_d) = 1 \text{ cm}$$




E. Bernreuther, F. Kahlhoefer, M. Kramer and P. Tunney, 1907.04346; S. Koren and R. McGehee, 1908.03559; Y. Hochberg, E. Kuflik, T. Volansky and J. G. Wacker, 1402.5143; Y. Hochberg, E. Kuflik and H. Murayama, 1512.07917

.....




LL, Y. Tsai, 1901.09936

$$m_{\eta_d} = 10, m_{\omega_d} = 12, \Lambda_d = 3 \text{ [GeV]}$$



Displaced Vertex Efficiency for (HL-)LHC


Current CMS:



Extended Higgs Sector

$$H \sim \mathbf{2}_{1/2,0}, H' \sim \mathbf{2}_{1/2,x_\Phi}, \Phi \sim \mathbf{1}_{0,x_\Phi}$$


$$V = -\frac{m_H^2}{2}h^2 + \frac{\lambda_H}{4}h^4 + \frac{m_{H'}^2}{2}h'^2 + \frac{\lambda_{H'}}{4}h'^4 - m_\Phi^2\varphi^2 + \lambda_\Phi\varphi^4 - \mu hh'\varphi$$



In large width and $f/m = 1$ limit

CMS: Results on a Light Z'

Efficiency drop due to soft muons (can't reach the muon chamber of $p_T < 3$ GeV)



Still probing an effective
mixing of 10^{-3} - 10^{-4}

Dark Pion Decays (ALP-Like)

Applies to arbitrary
flavor couplings

See also:

D. Aloni, Y. Soreq and M.
Williams, 1811.03474

A.1 $a \rightarrow \gamma\gamma$

A.2 $a \rightarrow \pi^+\pi^-\gamma$

A.3 $a \rightarrow \pi^+\pi^-\pi^0$

A.4 $a \rightarrow 3\pi^0$

A.5 $a \rightarrow \pi^0\pi^0\eta, \pi^+\pi^-$

A.6 $a \rightarrow \pi^0\pi^0\eta', \pi^+\pi^-$

A.7 $a \rightarrow \eta\eta\pi^0$


A.8 $a \rightarrow K^0\bar{K}^0\pi^0$

A.9 $a \rightarrow K^+K^-\pi^0$

A.10 $a \rightarrow K^+\bar{K}^0\pi^-, K^-K^0\pi^+$

A.11 $a \rightarrow \omega\omega, \phi\phi, K^{*+}K^{*-}, K^{*0}\bar{K}^{*0}$

A.12 $a \rightarrow \pi^+\pi^-\omega$



$$\Gamma_{(\hat{\pi} \rightarrow \mu^+ \mu^-)}^{-1} \simeq 45 \text{ cm} \left(\frac{f_a}{1 \text{ PeV}} \right)^2 \left(\frac{1 \text{ GeV}}{m_{\hat{\pi}}} \right) \sqrt{\frac{m_{\hat{\pi}}^2}{m_{\hat{\pi}}^2 - 4m_{\mu}^2}}$$

